

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# [4-(Methylsulfonyl)phenyl]acetic acid

Yan-Qin Yuan, Sheng-Rong Guo\* and Li-Jin Wang

Department of Chemistry, Lishui University, 323000 Lishui, Zhejiang Province, People's Republic of China

Correspondence e-mail: guosr9608@163.com


Received 23 October 2008; accepted 7 November 2008

 Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.048;  $wR$  factor = 0.140; data-to-parameter ratio = 12.7.

In the crystal structure of the title compound,  $\text{C}_9\text{H}_{10}\text{O}_4\text{S}$ , centrosymmetrically related molecules are linked into dimers by intermolecular  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds. Unconventional  $\text{C}-\text{H}\cdots\text{O}$  hydrogen-bond interactions are also present, connecting dimers into a three-dimensional network.

## Related literature

For general background on the properties of the title compound and its derivatives, see: Parimalan *et al.* (2008); Giridhar *et al.* (2006). For the crystal structures of related compounds, see: Guo & Yuan (2006); Hartung *et al.* (2004); Hodgson & Asplund (1991). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



## Experimental

### Crystal data

$\text{C}_9\text{H}_{10}\text{O}_4\text{S}$   
 $M_r = 214.23$   
 Monoclinic,  $P2_1/c$   
 $a = 19.086$  (7) Å  
 $b = 4.9711$  (18) Å  
 $c = 10.724$  (4) Å  
 $\beta = 106.102$  (6)°

$V = 977.5$  (6) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.32$  mm<sup>-1</sup>  
 $T = 298$  (2) K  
 $0.52 \times 0.30 \times 0.24$  mm

### Data collection

Bruker SMART APEX area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 2004)  
 $T_{\min} = 0.853$ ,  $T_{\max} = 0.928$

4462 measured reflections  
 1638 independent reflections  
 1502 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.019$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.140$   
 $S = 1.11$   
 1638 reflections

129 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.24$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.43$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O4}-\text{H4}\cdots\text{O3}^{\text{i}}$	0.82	1.87	2.693 (3)	175
$\text{C3}-\text{H3}\cdots\text{O2}^{\text{ii}}$	0.93	2.53	3.287 (3)	139
$\text{C1}-\text{H1B}\cdots\text{O1}^{\text{iii}}$	0.96	2.45	3.365 (4)	160

 Symmetry codes: (i)  $-x, -y + 2, -z + 1$ ; (ii)  $x, -y + \frac{3}{2}, z - \frac{1}{2}$ ; (iii)  $x, -y + \frac{3}{2}, z + \frac{1}{2}$ .

Data collection: SMART (Bruker, 2002); cell refinement: SAINT (Bruker, 2002); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

The authors thank Zhengjiang Natural Science Foundation for financial support (Y407240).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2263).

## References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555-1573.  
 Bruker (2002). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Giridhar, P., Vijaya Ramu, D. & Ravishankar, G. A. (2006). *Trop. Sci.* **43**, 92-95.  
 Guo, S.-R. & Yuan, Y.-Q. (2006). *Acta Cryst.* **E62**, o5707-o5708.  
 Hartung, J., Špehar, K., Svoboda, I. & Fuess, H. (2004). *Acta Cryst.* **E60**, o95-o96.  
 Hodgson, D. J. & Asplund, R. O. (1991). *Acta Cryst.* **C47**, 1986-1987.  
 Parimalan, R., Giridhar, P. & Ravishankar, G. A. (2008). *Ind. Crop. Prod.* **28**, 122-127.  
 Sheldrick, G. M. (2004). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.

## supporting information

*Acta Cryst.* (2008). E64, o2435 [doi:10.1107/S160053680803660X]

**[4-(Methylsulfonyl)phenyl]acetic acid**

Yan-Qin Yuan, Sheng-Rong Guo and Li-Jin Wang

**S1. Comment**

The title compound, C<sub>9</sub>H<sub>10</sub>O<sub>4</sub>S, and its substituted derivatives have been found to possess an auxin-like activity (Parimalan *et al.*, 2008; Giridhar *et al.*, 2006). These compounds, which are predominantly found in fruits, can be used in the synthesis of pharmaceutical intermediates, some perfumes and non-steroidal anti-inflammatory drugs.


In the molecule of the title compound (Fig. 1), bond lengths and angles agree well with those observed in similar compounds (Guo & Yuan, 2006; Hartung *et al.*, 2004; Hodgson & Asplund, 1991). The S=O bond lengths within the SO<sub>2</sub>Me group are not significantly different, with an average value of 1.4366 (10) Å. The average bond length for the two C—S bonds is 1.762 (3) Å. In the crystal packing, centrosymmetrically related molecules are linked into dimers by intermolecular O—H⋯O hydrogen bonds (Table 1) generating an eight-membered ring of graph set R<sup>2</sup><sub>2</sub>(8) (Bernstein *et al.*, 1995). The dimers are further linked into a three-dimensional network by unconventional C—H⋯O hydrogen bonding interactions. The corresponding phenylacetic acid derive without the SO<sub>2</sub>Me group (Hartung *et al.*, 2004) forms helical columns of single enantiomers linked by hydrogen bonds between the acidic proton of one molecule and the methoxy O atom of a neighbouring molecule, to give an overall racemic structure.

**S2. Experimental**

1-(4-Methanesulfonyl-phenyl)-ethanone (20 mmol), morpholine (60 mmol) and elemental sulfur (40 mmol) were added in a round-bottom flask and refluxed for 2 h at 398 K. A 3N solution of NaOH (20 ml) was then added, and the reaction mixture refluxed for an additional 30 min. After cooling, the mixture was filtered and the filtrate was acidified with HCl to pH 6. The solution was again filtered off and washed with ethyl acetate. The resulting aqueous fraction was finally acidified with diluted HCl, to yield the pure product as a white solid. Crystals suitable for X-ray analysis were obtained by slow evaporation of an ethanol/water (1:1 v/v) solution.

**S3. Refinement**

All H atoms were placed at calculated positions and constrained to ride on their parent atoms, with O—H = 0.82 Å, C—H = 0.93–0.97 Å and with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  or  $1.5 U_{\text{eq}}(\text{C}, \text{O})$  for hydroxy and methyl H atoms.

**Figure 1**

The molecular structure of the title compound, with the atom-labeling scheme and 50% probability displacement ellipsoids.

### [4-(Methylsulfonyl)phenyl]acetic acid

#### Crystal data

$C_9H_{10}O_4S$

$M_r = 214.23$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2ybc$

$a = 19.086\ (7)\ \text{\AA}$

$b = 4.9711\ (18)\ \text{\AA}$

$c = 10.724\ (4)\ \text{\AA}$

$\beta = 106.102\ (6)^\circ$

$V = 977.5\ (6)\ \text{\AA}^3$

$Z = 4$

$F(000) = 448$

$D_x = 1.456\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2824 reflections

$\theta = 2.2\text{--}27.7^\circ$

$\mu = 0.32\ \text{mm}^{-1}$

$T = 298\ \text{K}$

Block, colourless

$0.52 \times 0.30 \times 0.24\ \text{mm}$

#### Data collection

Bruker APEX area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 2004)

$T_{\min} = 0.853$ ,  $T_{\max} = 0.928$

4462 measured reflections

1638 independent reflections

1502 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.020$

$\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 1.1^\circ$

$h = -22 \rightarrow 22$

$k = -5 \rightarrow 5$

$l = -11 \rightarrow 12$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.048$

$wR(F^2) = 0.140$

$S = 1.11$

1638 reflections

129 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0848P)^2 + 0.3788P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.24\ \text{e \AA}^{-3}$

$\Delta\rho_{\min} = -0.43\ \text{e \AA}^{-3}$

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.38734 (3)	0.64558 (12)	1.01463 (5)	0.0408 (3)
O1	0.41630 (11)	0.5716 (4)	0.90951 (18)	0.0609 (6)
O2	0.36962 (11)	0.4358 (4)	1.09321 (19)	0.0575 (5)
O3	0.07977 (10)	0.9489 (5)	0.6087 (2)	0.0694 (7)
O4	0.00360 (10)	1.2830 (5)	0.6090 (2)	0.0677 (6)
H4	−0.0197	1.2097	0.5418	0.102*
C1	0.44823 (14)	0.8670 (5)	1.1176 (3)	0.0487 (6)
H1A	0.4947	0.7807	1.1501	0.073*
H1B	0.4296	0.9159	1.1889	0.073*
H1C	0.4537	1.0256	1.0701	0.073*
C2	0.30708 (12)	0.8358 (4)	0.9483 (2)	0.0361 (5)
C3	0.30404 (13)	1.0060 (5)	0.8448 (2)	0.0425 (6)
H3	0.3440	1.0226	0.8117	0.051*
C4	0.24133 (14)	1.1507 (5)	0.7913 (2)	0.0481 (6)
H4A	0.2393	1.2662	0.7221	0.058*
C5	0.18090 (13)	1.1260 (5)	0.8396 (2)	0.0434 (6)
C6	0.18554 (13)	0.9570 (6)	0.9437 (3)	0.0491 (6)
H6	0.1458	0.9413	0.9775	0.059*
C7	0.24791 (13)	0.8111 (5)	0.9986 (2)	0.0457 (6)
H7	0.2502	0.6975	1.0685	0.055*
C8	0.11162 (14)	1.2792 (6)	0.7781 (3)	0.0571 (7)
H8A	0.0838	1.2955	0.8410	0.069*
H8B	0.1244	1.4594	0.7574	0.069*
C9	0.06426 (13)	1.1525 (5)	0.6574 (3)	0.0451 (6)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0481 (4)	0.0392 (4)	0.0331 (4)	0.0129 (2)	0.0079 (3)	−0.0027 (2)
O1	0.0635 (11)	0.0752 (13)	0.0449 (11)	0.0256 (10)	0.0164 (9)	−0.0110 (10)
O2	0.0767 (13)	0.0387 (10)	0.0564 (12)	0.0120 (9)	0.0176 (10)	0.0082 (8)
O3	0.0530 (11)	0.0703 (14)	0.0648 (13)	0.0194 (10)	−0.0170 (9)	−0.0235 (11)
O4	0.0509 (11)	0.0665 (13)	0.0667 (14)	0.0203 (10)	−0.0155 (9)	−0.0117 (11)
C1	0.0470 (13)	0.0516 (15)	0.0415 (14)	0.0117 (11)	0.0024 (11)	−0.0031 (11)
C2	0.0381 (11)	0.0374 (12)	0.0294 (11)	0.0036 (9)	0.0038 (9)	−0.0038 (9)
C3	0.0420 (12)	0.0509 (14)	0.0338 (13)	0.0017 (10)	0.0090 (10)	0.0022 (10)

C4	0.0541 (14)	0.0492 (15)	0.0349 (13)	0.0037 (11)	0.0022 (11)	0.0066 (10)
C5	0.0387 (12)	0.0450 (14)	0.0378 (13)	0.0031 (10)	-0.0038 (10)	-0.0110 (10)
C6	0.0386 (12)	0.0585 (15)	0.0499 (15)	0.0000 (11)	0.0116 (11)	-0.0038 (13)
C7	0.0486 (13)	0.0489 (14)	0.0401 (14)	0.0013 (11)	0.0132 (11)	0.0062 (11)
C8	0.0487 (14)	0.0545 (16)	0.0557 (17)	0.0130 (13)	-0.0061 (12)	-0.0132 (13)
C9	0.0364 (12)	0.0471 (15)	0.0449 (14)	0.0044 (10)	-0.0001 (10)	0.0002 (11)

*Geometric parameters (Å, °)*

S1—O1	1.4342 (19)	C3—C4	1.378 (4)
S1—O2	1.439 (2)	C3—H3	0.9300
S1—C1	1.752 (3)	C4—C5	1.394 (4)
S1—C2	1.773 (2)	C4—H4A	0.9300
O3—C9	1.212 (3)	C5—C6	1.380 (4)
O4—C9	1.303 (3)	C5—C8	1.510 (3)
O4—H4	0.8200	C6—C7	1.379 (4)
C1—H1A	0.9600	C6—H6	0.9300
C1—H1B	0.9600	C7—H7	0.9300
C1—H1C	0.9600	C8—C9	1.497 (4)
C2—C3	1.384 (3)	C8—H8A	0.9700
C2—C7	1.386 (3)	C8—H8B	0.9700
O1—S1—O2	118.61 (12)	C3—C4—H4A	119.6
O1—S1—C1	108.87 (13)	C5—C4—H4A	119.6
O2—S1—C1	107.99 (13)	C6—C5—C4	118.7 (2)
O1—S1—C2	107.44 (11)	C6—C5—C8	120.8 (2)
O2—S1—C2	107.72 (12)	C4—C5—C8	120.5 (2)
C1—S1—C2	105.44 (11)	C7—C6—C5	121.3 (2)
C9—O4—H4	109.5	C7—C6—H6	119.3
S1—C1—H1A	109.5	C5—C6—H6	119.3
S1—C1—H1B	109.5	C6—C7—C2	119.2 (2)
H1A—C1—H1B	109.5	C6—C7—H7	120.4
S1—C1—H1C	109.5	C2—C7—H7	120.4
H1A—C1—H1C	109.5	C9—C8—C5	114.3 (2)
H1B—C1—H1C	109.5	C9—C8—H8A	108.7
C3—C2—C7	120.6 (2)	C5—C8—H8A	108.7
C3—C2—S1	119.18 (17)	C9—C8—H8B	108.7
C7—C2—S1	120.22 (18)	C5—C8—H8B	108.7
C4—C3—C2	119.5 (2)	H8A—C8—H8B	107.6
C4—C3—H3	120.3	O3—C9—O4	122.7 (2)
C2—C3—H3	120.3	O3—C9—C8	124.2 (2)
C3—C4—C5	120.8 (2)	O4—C9—C8	113.0 (2)
O1—S1—C2—C3	-35.3 (2)	C3—C4—C5—C8	-178.3 (2)
O2—S1—C2—C3	-164.15 (19)	C4—C5—C6—C7	-1.0 (4)
C1—S1—C2—C3	80.7 (2)	C8—C5—C6—C7	178.4 (2)
O1—S1—C2—C7	143.9 (2)	C5—C6—C7—C2	0.2 (4)
O2—S1—C2—C7	15.0 (2)	C3—C2—C7—C6	0.5 (4)

C1—S1—C2—C7	-100.1 (2)	S1—C2—C7—C6	-178.71 (19)
C7—C2—C3—C4	-0.3 (4)	C6—C5—C8—C9	-98.9 (3)
S1—C2—C3—C4	178.84 (19)	C4—C5—C8—C9	80.5 (3)
C2—C3—C4—C5	-0.4 (4)	C5—C8—C9—O3	-2.1 (4)
C3—C4—C5—C6	1.1 (4)	C5—C8—C9—O4	177.6 (3)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O4—H4 $\cdots$ O3 <sup>i</sup>	0.82	1.87	2.693 (3)	175
C3—H3 $\cdots$ O2 <sup>ii</sup>	0.93	2.53	3.287 (3)	139
C1—H1B $\cdots$ O1 <sup>iii</sup>	0.96	2.45	3.365 (4)	160

Symmetry codes: (i)  $-x, -y+2, -z+1$ ; (ii)  $x, -y+3/2, z-1/2$ ; (iii)  $x, -y+3/2, z+1/2$ .